

Higher l -adic Abel-Jacobi mappings and filtrations on Chow groups

Wayne Raskind*
 Dept. of Mathematics
 University of Southern California
 Los Angeles, CA 90089

Introduction

Let X be a smooth projective variety over a field k and denote by $CH^n(X)$ the group of codimension n cycles modulo rational equivalence on X . The structure of this group is largely unknown when $n \geq 2$, especially when k is a “large” field such as the complex numbers. Beginning with the work of Bloch [Bl3] and Beauville [Be1,2], the Chow ring of an abelian variety was found to have some structure arising from a filtration defined using Pontryagin products or the Fourier transform. In recent years, several people (Beilinson, Bloch, Murre, Nori, S. Saito, H. Saito; see [B1], [Bl4], [Mu], [N], [S1], [SH]) have made conjectures asserting the existence of certain filtrations on the Chow groups of any smooth projective variety. These conjectures would imply that the Chow groups have much more structure than has been hitherto uncovered. In this paper we define and study a filtration arising from the cycle map into l -adic étale cohomology. It is somewhat arithmetic in nature because it is necessary to first define it over a field which is finitely generated over its prime subfield. For varieties over larger fields, it may be defined by passing to the limit over finitely generated subfields. Others have studied

*Research supported by the National Science Foundation and the National Security Agency

this filtration, in particular Beauville (unpublished) and Jannsen (see [J4]). With their permission, I have included some of their results in this paper.

Each of the filtrations defined so far has its advantages and disadvantages. The conjectural filtration of Beilinson arising from a spectral sequence relating extensions of mixed motives and motivic cohomology would imply many outstanding and deep conjectures about algebraic cycles. We will not repeat this story in detail here since it is treated very clearly in the paper of Jannsen [J4]. The conjectural filtration of Murre is known to exist in some cases, and it would also imply some of these conjectures, but to establish its existence in general appears to require the resolution of some very difficult conjectures. The filtrations of H. Saito and S. Saito are defined in a very explicit way, but it seems difficult to do specific calculations. The filtration in this paper has the advantage that it may be “computed” in some cases, and the powerful tools of arithmetic algebraic geometry may be brought to bear on it. However, it has two basic disadvantages. First, it is not at all obvious that it is independent of l . Second, suppose X is defined over a field k which is finitely generated over its prime subfield and l is a prime number different from the characteristic of k . Jannsen [J1] has defined a cycle map into continuous étale cohomology (see §1):

$$c_n : CH^n(X) \otimes \mathbf{Q} \rightarrow H^{2n}(X, \mathbf{Q}_l(n)).$$

The following is a more general version of a conjecture first made by Soulé over finite fields and by Jannsen and the author over number fields:

Conjecture: c_n is injective.

This is known when $n=1$ (a consequence of the finite generation of $CH^1(X)$), but except for special classes of varieties, we have little information for $n \geq 2$.

The Hochschild-Serre spectral sequence:

$$H^r(k, H^q(\overline{X}, \mathbf{Q}_l(n))) \implies H^{r+s}(X, \mathbf{Q}_l(n))$$

defines a filtration on $H^{2n}(X, \mathbf{Q}_l(n))$, which we pull back to $CH^n(X) \otimes \mathbf{Q}$ via the cycle map c_n . We call this the *l-adic filtration* on $CH^n(X) \otimes \mathbf{Q}$. By definition, the kernel of the cycle map is contained in all the steps of the filtration. For this reason, until we have more information on this kernel, we

can only get results about the filtration modulo the kernel. We can define another filtration $\mathcal{F}^\bullet CH^n(X) \otimes \mathbf{Q}$ on the image of the cycle map by taking the intersection with the filtration obtained from the Hochschild-Serre spectral sequence. It is really this filtration about which we can say something.

In section 2 of this paper we refine the cycle map c_n to get *higher l -adic Abel-Jacobi maps* from certain parts of the Chow group to Galois cohomology groups. For example, suppose that $n = \dim X$ and let $A_0(X)$ denote the group of zero-cycles of degree zero modulo rational equivalence. Let $Alb(X)$ be the group of k -points of the Albanese variety of X and $T(X)$ the kernel of the Albanese map:

$$A_0(X) \rightarrow Alb(X).$$

Then we define a map:

$$d_n^2 : T(X) \otimes \mathbf{Q} \rightarrow H^2(k, H^{2n-2}(\overline{X}, \mathbf{Q}_l(n))).$$

This map generalizes the l -adic Abel-Jacobi map of Bloch [Bl1], which in this case is essentially the map obtained from Kummer theory on $Alb(X)$ (see §2 for more details).

For the results of this paper we will need to assume that if X is a smooth projective variety over a local field k with ring of integers $\mathcal{O}$, then X has a regular proper model over $\mathcal{O}$. Assuming this, the main result of this paper is:

Theorem 0.1: Let k be a function field in one variable over a finite field of characteristic $p \neq l$ and X a smooth projective variety of dimension n over k . Then the map d_n^2 is zero.

Since a global field is of cohomological dimension 2 for $\mathbf{Q}_l$ -cohomology, we have:

Corollary: For X as above of dimension n , we have $\mathcal{F}^i CH^n(X) \otimes \mathbf{Q} = 0$ for all $i \geq 2$.

The following remarks may help clarify the meaning of these results. First of all, Beilinson and Bloch have made the conjecture that if k is a global field and X is a smooth projective variety over k then the Albanese kernel $T(X)$

is a torsion group. Thus the theorem provides “evidence” for this conjecture for k a global field of positive characteristic by showing that the cycle classes of elements of $T(X)$ in $H^{2n}(X, \mathbf{Q}_l(n))$ are trivial. This should be true for k a number field as well, but we can only prove it when $H^{2n-2}(\overline{X}, \mathbf{Q}_l(n))$ is generated by the cohomology classes of algebraic cycles. In this case, Beauville showed me an easy proof which works over any field.

One of the key technical tools in this paper is the theory of continuous l -adic étale cohomology and the Hochschild-Serre spectral sequence relating the cohomology of a variety X over a field k with the cohomology over a separable closure $\overline{k}$ (see §1). Such spectral sequences do not exist in general for the usual l -adic cohomology defined by taking the inverse limit of cohomology with finite coefficients. Also, we rely on results of Jannsen on the Galois cohomology of global fields with coefficients in l -adic cohomology groups of algebraic varieties over $\overline{k}$ [J2]. Another key ingredient is Thomason’s purity theorem for $\mathbf{Q}_l$ -cohomology of schemes of finite type over the ring of integers in a local field of residue characteristic different from l [T]. This enables us to construct cycle maps over rings of integers of local fields.

The l -adic Abel-Jacobi maps d_n^i were discovered independently (and earlier) by Beauville and by Jannsen, who has also given a specific description of the map d_n^2 in terms of extension groups [J3].

At least for X smooth over a field, $CH^*(X) \otimes \mathbf{Q} \cong K_0(X) \otimes \mathbf{Q}$, and so the l -adic filtration may also be viewed as a filtration on $K_0(X) \otimes \mathbf{Q}$. There are Chern class maps:

$$K_j(X) \otimes \mathbf{Q} \rightarrow \bigoplus_{i,j} H^{2i-j}(X, \mathbf{Q}_l(i))$$

which one can hope are injective over an absolutely finitely generated field. These can be used to define similar filtrations on all K -groups. We hope to develop this line of investigation further in another paper.

I would like to thank U. Jannsen for explaining some of his unpublished results to me and allowing me to include some of them in this paper. J.-L. Colliot-Thélène listened to and helped me with several points in this paper, especially Proposition 3.2. A. Beauville informed me that one can easily

prove that the l -adic filtration is of length at most n on CH^n modulo the kernel of the cycle map. He also showed me a much simpler proof than the one I had originally proposed of the triviality of the higher Abel-Jacobi map in the case where the cohomology of $\overline{X}$ is generated by the classes of algebraic cycles. I am grateful to him for allowing me to include these here. The referee made several helpful comments which have improved the exposition. Finally, I would like to thank K. Paranjape and S. Saito for useful discussions.

During the long period of thinking about and writing this paper, I enjoyed the hospitality of the Max-Planck Institut für Mathematik in Bonn, Universität zu Köln, Universität Münster, Université de Paris-Sud and the CRM of the Universitat Autònoma de Barcelona. During my one-week visits to the aforementioned German universities, I was supported by the Deutsche Forschungsgemeinschaft.

1 Notation and Preliminaries

We denote by X a smooth, projective, geometrically connected variety over a field k and l a prime number different from $\text{char}.k$. Let $\overline{k}$ be a separable closure of k and $G = \text{Gal}(\overline{k}/k)$, $\overline{X} = X \times_k \overline{k}$. If k is a global field and S is a finite set of places of k , G_S denotes the Galois group of a maximal extension of k which is unramified outside S . For $\mathcal{X}$ a noetherian scheme, we denote by $CH^n(\mathcal{X})$ the Chow group of codimension n cycles modulo rational equivalence. For X proper and geometrically connected over a field, $CH_0(X)$ is the group of zero-cycles modulo rational equivalence and $A_0(X)$ the group of zero-cycles of degree zero modulo rational equivalence. The group of k -points of the Albanese variety of X will be denoted by $\text{Alb}(X)$ and $\alpha : A_0(X) \rightarrow \text{Alb}(X)$ denotes the Albanese map. We denote the kernel of this map by $T(X)$.

If M is an abelian group upon which G acts continuously, we denote the continuous Galois cohomology groups of G with values in M by $H^*(k, M)$ (see §1 for a definition of these groups). The subgroup of M consisting of elements killed by l^m is denoted by M_{l^m} and

$$T_l M = \varinjlim_m M_{l^m}.$$

We set $V_l M = T_l M \otimes_{\mathbf{Z}_l} \mathbf{Q}_l$. As usual, we denote by $\mathbf{Z}_l(1)$ the l -adic sheaf of l -primary roots of unity. For $n > 0$, $\mathbf{Z}_l(n)$ denotes $\mathbf{Z}_l(1)^{\otimes n}$; for $n < 0$, $\mathbf{Z}_l(n) = \text{Hom}(\mathbf{Z}_l(-n), \mathbf{Z}_l)$. We use similar notation for $\mathbf{Q}_l$ -sheaves.

Let X be a smooth variety over a field k and l a prime number different from the characteristic of k . Jannsen has defined a cycle map [J1]:

$$CH^n(X) \otimes \mathbf{Q} \rightarrow H^{2n}(X, \mathbf{Q}_l(n)),$$

where the group on the right is the continuous l -adic cohomology group with values in the l -adic sheaf $\mathbf{Q}_l(n)$. We briefly recall the definition of these groups. Let $(\mathcal{F}_m)$ be a projective system of sheaves of $\mathbf{Z}/l^m$ -modules on X . The functor which sends $(\mathcal{F}_m)$ to

$$\varprojlim_m H^0(X, (\mathcal{F}_m))$$

is left exact, and we define $H^i(X, (\mathcal{F}_m))$ to be the i -th right derived functor. When the inverse system is $\mathbf{Z}/l^m(j)$, we denote these groups by $H^i(X, \mathbf{Z}_l(j))$. Continuous étale cohomology with $\mathbf{Q}_l$ -coefficients is defined by tensoring the $\mathbf{Z}_l$ -cohomology with $\mathbf{Q}_l$.

We shall need the following properties of this cohomology which can be found in Jannsen's paper [J1].

(i) There is an exact sequence:

$$0 \rightarrow \varprojlim_m^1 H^{i-1}(X, (\mathcal{F}_m)) \rightarrow H^i(X, (\mathcal{F}_m)) \rightarrow \varprojlim_m H^i(X, \mathcal{F}_m) \rightarrow 0.$$

In particular, if $H^i(X, \mathbf{Z}/l^m(j))$ is finite for all i and m then

$$H^i(X, \mathbf{Z}_l(j)) \rightarrow \varprojlim_m H^i(X, \mathbf{Z}/l^m(j))$$

is an isomorphism. This is true if, for example, k is separably closed, finite or local. This is not true in general if k is a number field.

(ii) Let $\bar{k}$ be a separable closure of k , $G = \text{Gal}(\bar{k}/k)$ and $\bar{X} = X \times_k \bar{k}$. Then there is a Hochschild-Serre spectral sequence:

$$E_2^{p,q} = H^p(G, H^q(\bar{X}, \mathbf{Z}_l(j))) \implies H^{p+q}(X, \mathbf{Z}_l(j)).$$

Theorem 1.1 (Jannsen): Let X be a smooth projective variety over a field k . Then the Hochschild-Serre spectral sequence with $\mathbf{Q}_l$ -coefficients degenerates at E_2 .

Since this result has not been published, let us indicate the ingredients in the proof. The idea is to use Deligne's criterion for a spectral sequence to degenerate [D1]. In order to use this in this situation, one needs a good derived category of $\mathbf{Q}_l$ -sheaves. One cannot use Deligne's definition given in ([D2], 1.1.2d) because the cohomology groups with finite coefficients need not be finite. Jannsen has given a good definition of such a derived category in the general case. Once this is done one applies the Hard Lefschetz Theorem to satisfy Deligne's criterion. Ekedahl has published another definition of the derived category of $\mathbf{Q}_l$ -sheaves [E], and this should also suffice to prove the degeneration of the Hochschild-Serre spectral sequence.

2 The l -adic filtration

From the Hochschild-Serre spectral sequence for continuous cohomology:

$$E_2^{p,q} = H^p(k, H^q(\overline{X}, \mathbf{Q}_l(n))) \implies H^{p+q}(X, \mathbf{Q}_l(n)),$$

we get a filtration on $H^{2n}(X, \mathbf{Q}_l(n))$:

$$H^{2n}(X, \mathbf{Q}_l(n)) = F^0 \supset F^1 \supset \dots \supset F^{2n}$$

and $F^i/F^{i+1} = E_\infty^{i, 2n-i}$.

Definition 2.1: Let X be a smooth, projective, geometrically connected variety over a field k which is finitely generated over its prime subfield. Then we define:

$$F^i CH^n(X) \otimes \mathbf{Q} = c_n^{-1}[F^i H^{2n}(X, \mathbf{Q}_l(n))],$$

where c_n is the cycle map (see §1).

If k is not finitely generated, we define:

$$F^i CH^n(X) \otimes \mathbf{Q} = \lim_{\overrightarrow{L}} F^i CH^n(X_L) \otimes \mathbf{Q},$$

where L runs over all finitely generated fields contained in k for which X has a model over L .

Finally, set

$$\mathcal{F}^i CH^n(X) \otimes \mathbf{Q} = \text{Image } c_n \cap F^i H^{2n}(X, \mathbf{Q}_l(n)).$$

Of course, *a priori* $\mathcal{F}^\bullet CH^n(X) \otimes \mathbf{Q}$ is only a filtration on the image of the cycle map.

Let X be defined over a field L which is finitely generated over the prime subfield. By the definition of the filtration on $CH^n(X) \otimes \mathbf{Q}$ and the degeneration of the Hochschild-Serre spectral sequence (Theorem 1.1), we get maps:

$$d_n^i : \mathcal{F}^i CH^n(X) \otimes \mathbf{Q} \rightarrow H^i(k, H^{2n-i}(\overline{X}, \mathbf{Q}_l(n)))$$

which we call the *higher l -adic Abel-Jacobi mappings*. For X over arbitrary k , we define the higher Abel-Jacobi map as the limit of the maps defined above for X_L , where L runs over finitely generated subfields of k over which X has a model.

When $i=1$, these are the l -adic Abel-Jacobi mappings defined by Bloch [Bl1]. When $n = \dim X$, this is the Albanese map followed by the map given by Kummer theory for $Alb(X)$, as we now explain.

Let $A_0(X)$ be the subgroup of $CH^n(X)$ consisting of cycles of degree zero and let $Alb(X)$ denote the group of k -points of the Albanese variety of X . Consider the Albanese map

$$\alpha : A_0(X) \rightarrow Alb(X)$$

and its kernel $T(X)$. Then we have a commutative diagram with exact rows:

$$\begin{array}{ccccccc} 0 & \rightarrow & F^2 & \rightarrow & F^1 & \rightarrow & H^1(k, H^{2n-1}(X, \mathbf{Q}_l(n))) \rightarrow 0 \\ & & f \uparrow & & \uparrow & & \uparrow \\ 0 & \rightarrow & T(X) \otimes \mathbf{Q} & \rightarrow & A_0(X) \otimes \mathbf{Q} & \rightarrow & Alb(X) \otimes \mathbf{Q} \rightarrow 0 \end{array}.$$

Here the right vertical map is given as follows. By Poincaré duality and the Weil pairing, we have an isomorphism:

$$H^{2n-1}(\overline{X}, \mathbf{Q}_l(n)) \cong V_l(\text{Alb}(\overline{X})),$$

where V_l denotes the Tate $\mathbf{Q}_l$ -vector space of $\text{Alb}(\overline{X})$. Then the right vertical map is defined by doing Kummer theory modulo l^m on $\text{Alb}(X)$, passing to the inverse limit over m and tensoring with $\mathbf{Q}_l$. The commutativity of the right square is proved in the appendix to this paper. This induces the map f . Composing f with the map obtained from the spectral sequence:

$$F^2 \rightarrow E_{\infty}^{2, 2n-2},$$

we get a map

$$d_n^2 : T(X) \otimes \mathbf{Q} \rightarrow H^2(k, H^{2n-2}(\overline{X}, \mathbf{Q}_l(n))).$$

Remarks 2.1.1: (i) It would be interesting to find a complex analytic analogue of the maps d_n^i . Of course, d_n^1 may be defined in the complex category as the Abel-Jacobi map of Griffiths. The target group of our d_n^i may be reinterpreted as $\text{Ext}_{\mathbf{Q}_l}^i(\mathbf{Q}_l, H^{2n-i}(\overline{X}, \mathbf{Q}_l(n)))$ in the category of continuous Galois modules. The complex analogue of this is $\text{Ext}_{MHS}^i(\mathbf{Q}, H^{2n-i}(X, \mathbf{Q}))$, where MHS denotes the category of mixed Hodge structures. For $i = 1$, this works fine, as it is well-known that $\text{Ext}_{MHS}^1(\mathbf{Z}, H^{2n-1}(X, \mathbf{Z}))$ for $X/\mathbf{C}$ is the intermediate Jacobian of Griffiths. Unfortunately, all the higher Ext -groups are zero. Several people have suggested taking extensions in a suitable category of variations of Hodge structure.

(ii) The filtration F_B of S. Saito (see [Sa3], Definition (1-4)) is contained in the l -adic filtration. It is not clear to me whether this is the case for the filtration F_{BM} defined in (loc. cit. Definition (1-5)). For much more on the relations between various filtrations, see the paper of Jannsen [J4].

(iii) The l -adic Abel-Jacobi maps defined above have a conjectural mixed motivic interpretation. Recall that there are motivic cohomology groups with $\mathbf{Q}$ -coefficients $H_{\mathcal{M}}^i(X, \mathbf{Q}(j))$ with

$$H_{\mathcal{M}}^{2n}(X, \mathbf{Q}(n)) = CH^n(X) \otimes \mathbf{Q}.$$

These may be defined by:

$$H_{\mathcal{M}}^i(X, \mathbf{Q}(n)) = gr_{\gamma}^n K_{2n-i}(X) \otimes \mathbf{Q},$$

where γ denotes the gamma filtration in algebraic K -theory (here it is crucial that we are dealing with $\mathbf{Q}$ -coefficients because the integral theory is much less developed). Beilinson [B1] has made the conjecture that these groups have a canonical filtration coming from a spectral sequence:

$$Ext_{\mathcal{MM}}^p(\mathbf{Q}, H^q(X, \mathbf{Q}(j))) \implies H_{\mathcal{M}}^{p+q}(X, \mathbf{Q}(j)).$$

Here the subscript $\mathcal{MM}$ denotes the conjectural category of mixed motives on X and the Ext is taken in this category. The maps given above are in some sense “ l -adic realizations” of the corresponding maps obtained from this conjectural spectral sequence. This point of view is very useful for learning what to expect. In fact, if Beilinson’s conjecture on the existence of filtrations on Chow groups ([B1], [J4]) is true, then the filtration defined in this paper must agree with his (see [J4], Lemma 2.7).

We now prove some basic results on these filtrations.

Proposition 2.2 (Beauville): For X as above, we have

$$F^{n+j}CH^n(X) \otimes \mathbf{Q} = F^{n+1}CH^n(X) \otimes \mathbf{Q}$$

for all $j \geq 1$.

Proof: It suffices to show that the higher Abel-Jacobi maps d_n^i are zero for $i > n$. To prove this, let d be the dimension of X and let Y be a smooth hyperplane section of X . Then the diagram:

$$\begin{array}{ccc} F^{n+j}CH^n(X) \otimes \mathbf{Q} & \rightarrow & H^{n+j}(k, H^{n-j}(\overline{X}, \mathbf{Q}_l(n))) \\ \downarrow & & \downarrow \\ F^{n+j}CH^{d+j}(X) \otimes \mathbf{Q} & \rightarrow & H^{n+j}(k, H^{2d-n+j}(\overline{X}, \mathbf{Q}_l(d+j))) \end{array}$$

is commutative. Here the left vertical arrow is given by intersecting cycles on X with the class of the $(d+j-n)$ -fold intersection of Y with itself, and the right vertical arrow is given by cupping $(d+j-n)$ -times with the class of Y in $H^2(X, \mathbf{Q}_l(1))$. This commutativity is easily seen from the compatibility of the Hochschild-Serre spectral sequence and the cycle map with intersection products ([J1], Lemma 6.14). Now since $j \geq 1$, the bottom left group is zero. But the right vertical arrow is an isomorphism by the Hard Lefschetz Theorem, and hence the image of the top horizontal arrow is trivial. This

proves the proposition.

Recall that the cohomology group $H^i(\overline{X}, \mathbf{Q}_l)$ is said to be *algebraic* if it is generated by Tate twists of cohomology classes of algebraic cycles. Thus if i is odd, this means that the group is zero, and if $i = 2j$ is even then the cycle map:

$$CH^j(\overline{X}) \otimes \mathbf{Q}_l \rightarrow H^{2j}(\overline{X}, \mathbf{Q}_l(j))$$

is surjective (we keep the Tate twists because we will need them later). A basic conjecture of Bloch ([Bl2], Conjecture 0.4 in the case of surfaces) predicts that if X is defined over $\mathbf{C}$ and the cohomology of X is supported on codimension 1 subschemes, then the Albanese map:

$$\alpha : A_0(X) \rightarrow Alb(X)$$

is an isomorphism. We now give an argument of Beauville to show how a weaker form of this conjecture is implied by the conjecture on injectivity of the cycle map for varieties over finitely generated fields (see the introduction to this paper).

Proposition 2.3 (Beauville): Let X be a smooth projective variety of dimension d over a field k . Let i be a positive integer and assume that the cohomology groups $H^{2n-2i}(\overline{X}, \mathbf{Q}_l(i))$ and $H^{2d-2n+2i}(\overline{X}, \mathbf{Q}_l(d-n+i))$ are generated by algebraic cycles. Then the map:

$$F^{2i}CH^n(X) \otimes \mathbf{Q} \rightarrow H^{2i}(k, H^{2n-2i}(\overline{X}, \mathbf{Q}_l(n)))$$

is zero.

Proof: First note that for any finite extension L/k , the natural map:

$$H^{2i}(k, H^{2n-2i}(\overline{X}, \mathbf{Q}_l(n))) \rightarrow H^{2i}(L, H^{2n-2i}(\overline{X}, \mathbf{Q}_l(n)))$$

is injective because its kernel is killed by $[L : k]$. By the algebraicity assumption, there is a finite extension L/k over which $G = Gal(\overline{k}/L)$ acts trivially on $H^{2n-2i}(\overline{X}, \mathbf{Q}_l(n-i))$ and on $H^{2d-2n+2i}(\overline{X}, \mathbf{Q}_l(d-n+i))$. Thus we may and do assume that $Gal(\overline{k}/k)$ acts trivially on these groups. Let $Y_1, \dots, Y_\nu$ be codimension $d-n+i$ cycles on X whose cohomology classes $\xi_1, \dots, \xi_\nu$

form a basis for $H^{2d-2n+2i}(\overline{X}, \mathbf{Q}_l(d-n+i))$. Then the following diagram is commutative:

$$\begin{array}{ccc} F^{2i}CH^n(X) \otimes \mathbf{Q} & \rightarrow & H^{2i}(k, H^{2n-2i}(\overline{X}, \mathbf{Q}_l(n))) \\ \downarrow & & \downarrow \\ [F^{2i}CH^{d+i}(X) \otimes \mathbf{Q}]^\nu & \rightarrow & [H^{2i}(k, H^{2d}(\overline{X}, \mathbf{Q}_l(d+i)))]^\nu \end{array}.$$

Here the left vertical arrow is given by intersecting with the class of Y_j on the j -th factor. The right vertical arrow is given by cupping with the class of ξ_j on the j -th factor; it is an isomorphism by the assumption that the cohomology is algebraic and Galois acts trivially on $H^{2n-2i}(\overline{X}, \mathbf{Q}_l(n-i))$. The commutativity is easily established by the compatibility of the Hochschild-Serre spectral sequence and the cycle map with intersection product. Now the bottom left group is zero if $i \geq 1$. Hence the top horizontal map is zero. This completes the proof of the proposition.

Corollary 2.4: Let X be a smooth projective variety of dimension d over an algebraically closed field k . assume that for $d \leq i \leq 2d-2$, the cohomology group $H^i(\overline{X}, \mathbf{Q}_l)$ is algebraic. Let K be a finitely generated field of definition of X and assume that the cycle map:

$$c_d : CH^d(X_M) \otimes \mathbf{Q} \rightarrow H^{2d}(X_M, \mathbf{Q}_l(d))$$

is injective over any finitely generated field M containing K . Then the Albanese map:

$$\alpha : A_0(X) \rightarrow Alb(X)$$

is an isomorphism. In particular, if $k = \mathbf{C}$ and X is a surface with $H^0(X, \Omega_X^2) = 0$, then injectivity of the cycle map c_2 over all finitely generated fields containing K implies Bloch's conjecture.

Proof: Let a be in the kernel of α . Then a is defined over some finitely generated field M . The assumption on injectivity of the cycle map implies that the maps:

$$F^iCH^d(X_M) \otimes \mathbf{Q} \rightarrow H^i(M, H^{2d-i}(\overline{X}, \mathbf{Q}_l(d)))$$

are injective. If $i \geq 2$ then the assumption that the cohomology is algebraic and Proposition 2.3 imply that the map is zero. When $i = 1$, the map may

be identified with α over M followed by Kummer theory for $\text{Alb}(X_M)$ (see the appendix to this paper). Hence α is injective up to torsion. But by Roitman's theorem [Ro], [M2], the kernel of α has no torsion, so it is zero. The last statement follows immediately from the exponential sequence, Serre duality and the comparison theorem between singular and étale cohomology.

Remark 2.4.1: This result should be true under the weaker hypothesis that for each $d \leq i \leq 2d - 2$, the cohomology group $H^i(\overline{X}, \mathbf{Q}_l)$ is supported on codimension one subschemes. This is Bloch's conjecture in the general case.

Proposition 2.5: Let X be a smooth, projective variety over a field k which is finitely generated over its prime subfield and l a prime number different from the characteristic of k . Then the image of the cycle map:

$$c_n : CH^n(X) \otimes \mathbf{Q}_l \rightarrow H^{2n}(X, \mathbf{Q}_l(n))$$

is a finite dimensional $\mathbf{Q}_l$ -vector space.

Proof: There exist an open subset U of $\text{Spec} \mathbf{Z}[1/l]$, a regular ring A of finite type and smooth over U with fraction field k , and a smooth proper scheme $\mathcal{X}$ over A with generic fibre X . There is a cycle map:

$$CH^n(\mathcal{X}) \otimes \mathbf{Q}_l \rightarrow H^{2n}(\mathcal{X}, \mathbf{Q}_l(n))$$

(see e.g. ([S2], Lemma 5.3) or Proposition 2.9 below). Now $H^{2n}(\mathcal{X}, \mathbf{Q}_l(n))$ is a finite dimensional $\mathbf{Q}_l$ vector space (see e.g. [CT-R], Proof of Théorème 1.1, (d)) and since the map

$$CH^n(\mathcal{X}) \otimes \mathbf{Q}_l \rightarrow CH^n(X) \otimes \mathbf{Q}_l$$

is surjective, we get the proposition.

Corollary 2.6: With hypotheses as in Proposition 2.5, for any nonnegative integer i , the image of the higher l -adic Abel-Jacobi map:

$$F^i CH^n(X) \otimes \mathbf{Q}_l \rightarrow H^i(k, H^{2n-i}(\overline{X}, \mathbf{Q}_l(n)))$$

is a finite dimensional $\mathbf{Q}_l$ vector space.

Proposition 2.5 and Corollary 2.6 are not very refined results since their proofs only use the existence of a smooth proper model of X over an open subset of a model of k over an open subset of $\text{Spec}\mathbf{Z}$. In order to get finer results, we need to assume the existence of a regular proper model $\mathcal{Y}$ of k over $\mathbf{Z}$ together with a regular proper model of X over $\mathcal{Y}$. In the rest of this section we assume we have such models. Our philosophy is that the image of the higher Abel-Jacobi map d_n^i lies in the *unramified part* of $H^i(k, H^{2n-i}(\overline{X}, \mathbf{Q}_l(n)))$, in the following sense.

Definition 2.7: Let k be a field which is finitely generated over its prime subfield, l a prime number which is invertible in k and $\mathcal{Y}$ a regular proper model of k over $\mathbf{Z}$. Let F be an l -adic sheaf on $\mathcal{Y}$. We denote by $H_{nr}^i(k, F)^{(l)}$ the $\mathbf{Q}_l$ -vector space:

$$\text{Ker}[H^i(k, F) \rightarrow \prod_{y \in \mathcal{Y}^{1'}} H_y^{i+1}(A_y, F)].$$

Here $\mathcal{Y}^{1'}$ is the set of codimension one points of $\mathcal{Y}$ which do not lie above l and A_y is the valuation ring of y . By abuse of notation, we also denote by F the sheaves on k and A_y induced by F . If F is a locally constant sheaf and l is invertible on $\mathcal{Y}$, it follows easily from purity that this is the usual definition of unramified cohomology (see [CT-O], for example).

Now let X be a smooth projective variety over k . Assume that we have a regular proper scheme $\mathcal{Y}$ over $\mathbf{Z}[1/l]$ with function field k and a regular proper scheme $\mathcal{X}$ together with a morphism:

$$f : \mathcal{X} \rightarrow \mathcal{Y}$$

whose generic fibre is X/k . We consider the l -adic sheaf $F = R^{2n-i}f_*\mathbf{Q}_l(n)$, which is locally constant on the smooth locus of f , and we can define the unramified cohomology of k with values in this sheaf. Then we have:

Conjecture 2.8: With notation as above, the image of the higher Abel-Jacobi map d_n^i is contained in $H_{nr}^i(k, F)^{(l)}$.

Remark 2.8.1: In the definition of unramified cohomology and the conjecture, we have ignored points of $\mathcal{Y}$ lying above l , not because we don't think

these are important, but rather because we are unable to formulate an intelligent conjecture on the image of the Abel-Jacobi map at these places. What is needed is a higher dimensional analogue of the finite part, exponential part and geometric part of Bloch-Kato [BK]. This will involve l -adic cohomology over a complete discretely valued field with nonperfect residue field of characteristic l , and this is not very well developed at the moment.

Another way to formulate Conjecture 2.8 is that the filtration on the Chow groups defined in 2.1 should extend to $\mathcal{X}$. We define this filtration using the following result:

Proposition 2.9: Let R be a ring which is of type (i), (ii) or (iii) below:

- (i) of finite type over $\mathbf{Z}$
- (ii) a localization of a ring as in (i)
- (iii) the henselization of a ring as in (ii)

Let $\mathcal{X}$ be a regular scheme of finite type over R and assume that l is invertible on $\mathcal{X}$. Then there is a cycle map:

$$c_n : CH^n(\mathcal{X}) \rightarrow H^{2n}(\mathcal{X}, \mathbf{Q}_l(n)).$$

Proof: Thomason [T] has proved a purity theorem for $\mathbf{Q}_l$ -cohomology of such schemes, and then the existence of a cycle map may be proven by copying the proof in e.g. ([M1], Chapter VI, §6).

Definition 2.10: Let $f : \mathcal{X} \rightarrow \mathcal{Y}$ be a proper morphism of schemes of finite type over a ring R as in 2.9 and l a prime number invertible on $\mathcal{Y}$. Then we have the Leray spectral sequence for continuous cohomology ([J1], 3.10):

$$H^p(\mathcal{Y}, R^q f_* \mathbf{Q}_l(n)) \implies H^{p+q}(\mathcal{X}, \mathbf{Q}_l(n)).$$

This defines a filtration on $H^{2n}(\mathcal{X}, \mathbf{Q}_l(n))$ which we can pull back to $CH^n(\mathcal{X}) \otimes \mathbf{Q}$ via c_n . We denote the resulting filtration $F^\bullet CH^n(\mathcal{X}) \otimes \mathbf{Q}$. Of course, this filtration may depend on $\mathcal{Y}$, but this will be fixed in what follows.

The following is a properness conjecture for the filtration on a regular proper scheme over a discrete valuation ring.

Conjecture 2.11: Let A be a discrete valuation ring as in 2.9(ii) or (iii) with fraction field k and l a prime number invertible on A . Let $\mathcal{X}$ be a regular proper scheme of finite type over A with generic fibre X . Then the natural map:

$$F^iCH^n(\mathcal{X}) \otimes \mathbf{Q} \rightarrow F^iCH^n(X) \otimes \mathbf{Q}$$

is surjective.

Proposition 2.12: With notation as in 2.11, assume that A is Henselian and $\mathcal{X}$ is smooth over A . Then Conjecture 2.11 is true for $\mathcal{X}$.

Proof: We need a lemma.

Lemma 2.13: Let A be a Henselian discrete valuation ring with fraction field k and (F_m) a projective system of locally constant l -primary sheaves on A . Then the natural map on continuous cohomology groups:

$$H^i(A, (F_m)) \rightarrow H^i(k, (F_m))$$

is injective.

Proof: Let s be the closed point of $S = \text{Spec}A$. For each m we have an exact sequence:

$$(*) \dots \rightarrow H^{i-1}(k, F_m) \xrightarrow{f} H_s^i(A, F_m) \rightarrow H^i(A, F_m) \rightarrow H^i(k, F_m) \rightarrow \dots$$

Now the second group in this sequence is isomorphic to $H^{i-2}(s, F_m(-1))$ by purity for locally constant sheaves on A . The map f is split surjective with a splitting being given by the map (π a uniformizing parameter of k):

$$H^{i-2}(s, F_m(-1)) \cong H^{i-2}(A, F_m(-1)) \xrightarrow{\cup \pi} H^{i-1}(k, F_m).$$

Hence the map obtained from f above by passing to the inverse limit over m is surjective as well. A similar statement holds for the case of the map

$$\varprojlim_m^1 H^{i-1}(k, F_m) \rightarrow \varprojlim_m^1 H^{i-2}(s, F_m(-1)).$$

Then by the property (i) of continuous cohomology discussed in §1 above, the map:

$$H^{i-1}(k, (F_m)) \rightarrow H^{i-2}(s, (F_m))$$

is surjective. Now there is a localization sequence like (*) above for continuous cohomology with support and a purity theorem for the cohomology with support in s ([J1], 3.8 and same argument as in Theorem 3.17 of that paper). We conclude that the map

$$H^i(A, (F_m)) \rightarrow H^i(k, (F_m))$$

is injective. This completes the proof of the lemma.

Returning to the proof of the proposition, since $\mathcal{X}$ is smooth over A , the sheaves $F_{i,m} := R^i f_* \mathbf{Z}/l^m(n)$ are locally constant on A (smooth and proper base change theorem), so the maps

$$H^j(A, (F_{i,m})) \rightarrow H^j(k, (F_{i,m}))$$

are injective. Then the Leray spectral sequence

$$H^p(A, R^q f_*(F_m)) \Longrightarrow H^{p+q}(\mathcal{X}, (F_m))$$

degenerates at E_2 because it does so for X/k (Theorem 1.1). Now an easy induction starting from the fact that the map

$$CH^n(\mathcal{X}) \rightarrow CH^n(X)$$

is surjective proves the proposition.

Remarks 2.14: (i) When $\mathcal{X}$ is not smooth over A , it seems much more difficult to prove properness of the filtration. In fact, it is not even known if the map

$$F^1 CH^n(\mathcal{X}) \otimes \mathbf{Q} \rightarrow F^1 CH^n(X) \otimes \mathbf{Q}$$

is surjective. In Proposition 3.2 below we prove this when $n = \dim X$, A is Henselian and k with finite or separably closed residue field.

(ii) It should be possible to prove Proposition 2.12 for any discrete valuation ring A using results of Gillet [Gi]. His method may be used to show that if F is a locally constant sheaf on A with finite stalks, then the natural map:

$$H^i(A, F) \rightarrow H^i(k, F)$$

is injective. We need this for continuous cohomology. In some sense, our result for Henselian rings is not what we really want because the fraction field k is usually not finitely generated over the prime field. However, as will be shown in §3, in practice the Henselian case is very helpful.

3 Reductions in the proof of the Main Theorem

In this section we show how to reduce Theorem 0.1 to a Hasse principle for Galois cohomology which has been proved in some cases by Jannsen. In order to do this, we need some local considerations and, unfortunately, a hypothesis:

Hypothesis 3.1: Let A be a discrete valuation ring with fraction field k and X a smooth projective variety over k .

(Resolution of singularities) X has a regular proper model $\mathcal{X}$ over A .

Hypothesis 3.1 will be needed in order to have a cycle map for $\mathcal{X}$ (see Proposition 2.9) and for Proposition 3.2 below.

The following result may be found in ([B1], Lemma 2.2.6(b)). We give some of the details of the proof outlined there as well as the slight generalization we require.

Proposition 3.2: Let A be a Henselian discrete valuation ring with fraction field k and residue field $\mathbf{F}$ which is of characteristic different from l . Assume that $\mathbf{F}$ is either finite or separably closed. Let X be a smooth, projective, geometrically connected variety of dimension n over k and ξ an element of $CH^n(X) \otimes \mathbf{Q}$ of degree 0. Then there exists a finite extension L/k with valuation ring B , a regular proper model $\mathcal{X}$ of X_L over B and a cycle in $CH^n(\mathcal{X}) \otimes \mathbf{Q}_l$ which restricts to ξ_L in $CH^n(X_L) \otimes \mathbf{Q}_l$ and whose cohomology class in $H^{2n}(\mathcal{X}, \mathbf{Q}_l(n))$ is zero.

Proof: If ξ is a zero-cycle of degree zero on X , its cohomology class is trivial in $H^{2n}(\overline{X}, \mathbf{Q}_l(n))$. We claim that its cohomology class is zero in $H^{2n}(X, \mathbf{Q}_l(n))$. To see this, note that by Lemma 3.3 below, the group $A_0(X)$ is an extension of a torsion group by a group which is divisible prime to the residue characteristic of k . Now the group $H^{2n}(X, \mathbf{Z}_l(n))$ contains no nontrivial l -divisible subgroup ([J1], Cor. 4.9). Thus the restriction of the integral cycle map to cycles of degree zero:

$$A_0(X) \rightarrow H^{2n}(X, \mathbf{Z}_l(n))$$

has torsion image, so the class of ξ in $H^{2n}(X, \mathbf{Q}_l(n))$ is trivial, as claimed.

Assume for the moment that X is a curve. Let $\mathcal{X}$ be a regular proper model of X over A with special fibre Y . There is a commutative diagram with exact top row:

$$\begin{array}{ccccccc} CH^0(Y) \otimes \mathbf{Q}_l & \rightarrow & CH^1(\mathcal{X}) \otimes \mathbf{Q}_l & \rightarrow & CH^1(X) \otimes \mathbf{Q}_l & \rightarrow & 0 \\ & & \downarrow & & \downarrow & & \\ & & H^2(\mathcal{X}, \mathbf{Q}_l(1)) & \rightarrow & H^2(X, \mathbf{Q}_l(1)) & & \end{array}.$$

The vertical maps are cycle maps and the top sequence is the exact sequence for the Chow groups of a scheme, an open subscheme and its complement. By Lemma 3.4 below, we can identify $H^2(\mathcal{X}, \mathbf{Q}_l(1))$ with $(CH^0(Y_{\overline{\mathbf{F}}}) \otimes \mathbf{Q}_l)^{*Gal(\overline{\mathbf{F}}/\mathbf{F})}$, and the cycle map

$$CH^1(\mathcal{X}) \otimes \mathbf{Q} \rightarrow H^2(\mathcal{X}, \mathbf{Q}_l(1))$$

constructed in Lemma 2.9 is given by intersecting with components of the geometric special fibre. Here $*$ denotes $\mathbf{Q}_l$ -linear dual. Now the intersection

pairing on $CH^0(Y)/[Y]$ in $\mathcal{X}$ is nondegenerate, where $[Y]$ is the class of the whole special fibre ([L], Ch. 12, §4), and it induces an isomorphism:

$$CH^0(Y)/[Y] \otimes \mathbf{Q}_l \rightarrow [CH^0(Y_{\overline{\mathbf{F}}})^0 \otimes \mathbf{Q}_l]^{\ast Gal(\overline{\mathbf{F}}/\mathbf{F})},$$

where the superscript 0 means the kernel of the degree map.

If ξ is a zero-cycle of degree zero on X and $\tilde{\xi}$ is any lifting of ξ to a codimension 1 cycle on $\mathcal{X}$, then the class of $\tilde{\xi}$ in $CH^0(Y)$ is of total degree zero, where total degree means the sum of the degrees on each irreducible component of Y . Hence we can modify $\tilde{\xi}$ by an element of $CH^0(Y) \otimes \mathbf{Q}_l$ so that its class in $[CH^0(Y_{\overline{\mathbf{F}}}) \otimes \mathbf{Q}_l]^{\ast}$ is trivial. This proves the lemma in the case of a curve. For the higher dimensional case, let ξ be an element of $CH^n(X) \otimes \mathbf{Q}$ of degree zero. Then passing to a finite extension of k if necessary (if k is not perfect), we may find a smooth, projective, geometrically connected curve C passing through ξ ([C], [AK]). Let B be the integral closure of A in L and let $\mathcal{C}$ be a regular proper model of C over B . Then by blowing up, the rational map $\mathcal{C} \rightarrow \mathcal{X}$ may be resolved to get a morphism:

$$f : \tilde{\mathcal{C}} \rightarrow \mathcal{X},$$

with $\tilde{\mathcal{C}}$ regular.

Let Z, Y be the special fibres of $\tilde{\mathcal{C}}$ and $\mathcal{X}$, respectively. Then the following diagram commutes:

$$\begin{array}{ccc} CH^1(\tilde{\mathcal{C}}) \otimes \mathbf{Q}_l & \rightarrow & CH^n(\mathcal{X}) \otimes \mathbf{Q}_l \\ \downarrow & & \downarrow \\ [CH^0(Z_{\overline{\mathbf{F}}}) \otimes \mathbf{Q}_l]^{\ast} & \rightarrow & [CH^0(Y_{\overline{\mathbf{F}}}) \otimes \mathbf{Q}_l]^{\ast} \end{array}.$$

Here the top horizontal map is the push-forward map and the bottom horizontal map is defined as the dual of the pullback map on irreducible components.

By the first part of the proof, we can find a cycle $\tilde{\xi}$ in $CH^1(\tilde{\mathcal{C}}) \otimes \mathbf{Q}_l$ whose image in $CH^1(C) \otimes \mathbf{Q}_l$ is equal to ξ and whose class is trivial in $CH^0(Z) \otimes \mathbf{Q}_l$. Pushing $\tilde{\xi}$ to $CH^n(\mathcal{X}) \otimes \mathbf{Q}_l$, we get our desired cycle. This completes the proof of the proposition.

We now prove the two lemmas which were used in the proof of Proposition 3.2.

Lemma 3.3: Let k be a Henselian discrete valuation field. Assume that the residue field $\mathbf{F}$ is finite or separably closed. Let X be a smooth projective variety over k . Then the group $A_0(X)$ is an extension of a torsion group by a group which is divisible prime to the characteristic of $\mathbf{F}$.

Proof: As in the latter part of the proof of Proposition 3.2, given any zero-cycle on X , after passing to a finite extension of k if necessary, we may find a smooth, projective, geometrically connected curve going through its support. By an easy norm argument, we see that it suffices to prove the result for the case of curves. This is implied by the following fact applied to the Jacobian of the curve:

Fact: Let k be as in the hypothesis of Lemma 3.3 and A an abelian variety over k . Then the group $A(k)$ is an extension of a finite group by a group which is divisible prime to the characteristic of $\mathbf{F}$.

This is well-known if the residue field is finite. To see it when the residue field is separably closed, let $\mathcal{A}$ be the Néron model of A over the valuation ring R of k with special fibre $\tilde{\mathcal{A}}$ (see [BLR], Ch. 1, §1.3, Cor. 2 for the existence of $\mathcal{A}$). Let $A_0(k)$ be the subgroup of $A(k)$ consisting of points which specialize to elements in the connected component of identity $\tilde{\mathcal{A}}_0$ of $\tilde{\mathcal{A}}$. Then the quotient $A(k)/A_0(k)$ is finite. The group $\tilde{\mathcal{A}}_0(\mathbf{F})$ is the group of points of a connected algebraic group over the separably closed field $\mathbf{F}$, and hence is divisible prime to the characteristic of $\mathbf{F}$. Let $A_1(k)$ be the kernel of the surjective reduction map:

$$A_0(k) \rightarrow \tilde{\mathcal{A}}_0(\mathbf{F}).$$

Then $A_1(k)$ is a pro- p group, where p is the characteristic of $\mathbf{F}$. Hence it is uniquely divisible prime to p . Putting everything together, we get the fact.

Lemma 3.4: Let A be a Henselian local ring with residue field $\mathbf{F}$ which is finite or separably closed. Let $\mathcal{X}$ a regular proper flat integral scheme of dimension n over A with special fibre Y . Let l be a prime number different from the characteristic of $\mathbf{F}$. Then

$$(\dagger) \ H^{2n}(\mathcal{X}, \mathbf{Q}_l(n)) \cong [CH^0(Y_{\overline{\mathbf{F}}}) \otimes \mathbf{Q}_l]^*{}^{Gal(\overline{\mathbf{F}}/\mathbf{F})},$$

where $*$ denotes $\mathbf{Q}_l$ -dual. The map

$$CH^n(\mathcal{X}) \rightarrow [CH^0(Y_{\overline{\mathbf{F}}}) \otimes \mathbf{Q}_l]^*$$

obtained by composing this isomorphism with the cycle map in Proposition 2.9 is given by intersecting with components of the geometric special fibre.

Proof: This is known, but I could not find a good reference. By the proper base change theorem, we have:

$$H^{2n}(\mathcal{X}, \mathbf{Q}_l(n)) = H^{2n}(Y, \mathbf{Q}_l(n)).$$

This last group may be easily calculated as follows. Since the étale cohomology with $\mathbf{Q}_l$ -coefficients of Y and Y_{red} is the same, we may assume that Y is reduced. Let Y_{sing} be the singular locus of Y . Then there is an exact sequence for cohomology with compact support:

$$\dots \rightarrow H_c^{2n-1}(Y_{sing}, \mathbf{Q}_l(n)) \rightarrow H_c^{2n}(Y - Y_{sing}, \mathbf{Q}_l(n)) \rightarrow H_c^{2n}(Y, \mathbf{Q}_l(n)) \rightarrow H_c^{2n}(Y_{sing}, \mathbf{Q}_l(n)).$$

First assume that $\mathbf{F}$ is separably closed. Then since Y_{sing} is a proper closed subset of Y , the groups on the ends are zero. By Poincaré duality for $Y - Y_{sing}$, we have:

$$H_c^{2n}(Y - Y_{sing}, \mathbf{Q}_l(n)) \cong H^0(Y - Y_{sing}, \mathbf{Q}_l)^*,$$

and this last group is easily seen to be isomorphic to the $\mathbf{Q}_l$ -dual of $CH^0(Y) \otimes \mathbf{Q}_l$. If $\mathbf{F}$ is finite, then we have an exact sequence:

$$0 \rightarrow H^1(\mathbf{F}, H^{2n-1}(Y_{\overline{\mathbf{F}}}, \mathbf{Q}_l(n))) \rightarrow H^{2n}(Y, \mathbf{Q}_l(n)) \rightarrow H^{2n}(Y_{\overline{\mathbf{F}}}, \mathbf{Q}_l(n))^{Gal(\overline{\mathbf{F}}/\mathbf{F})} \rightarrow 0.$$

By Deligne's theorem ([D2], Theorem 3.3.1), the weights of the geometric Frobenius acting on $H^{2n-1}(Y_{\overline{\mathbf{F}}}, \mathbf{Q}_l(n))$ are ≤ -1 and hence are nonzero. This implies that the group on the left is zero. Using what we have just proved for separably closed residue fields, we get the identification (†). We leave the rest to the reader.

Remarks 3.5: (i) If the field k is perfect, then there is no need to pass to a finite extension in Proposition 3.2.

(ii) It is expected that, at least after tensoring with $\mathbf{Q}$, a cycle of arbitrary codimension which is homologically equivalent to zero on the generic fibre X can be lifted to one which is homologically equivalent to zero on any regular proper model $\mathcal{X}$.

Proposition 3.6: Let k be a function field in one variable over a finite field, X a smooth, projective variety of dimension n over k and l a prime number different from char. k . Put

$$M = H^{2n-2}(\overline{X}, \mathbf{Q}(n)).$$

Let S be a finite set of places of k including the bad reduction places of X . Then if z is a zero-cycle in the Albanese kernel $T(X) \otimes \mathbf{Q}$, $d_n^2(z)$ is contained in the image of

$$\text{Ker}[H^2(G_S, M) \rightarrow \bigoplus_{v \in S} H^2(k_v, M)]$$

in $H^2(k, M)$, where d_n^2 is the map defined in Section 2.

Proof: Let $U = \text{Spec } \mathcal{O}_{k,S}$. Then X extends to a smooth, proper scheme $\mathcal{X}_U$ over U and we can define a cycle map:

$$CH^n(\mathcal{X}_U) \rightarrow H^{2n}(\mathcal{X}_U, \mathbf{Q}_l(n)).$$

Consider the Leray spectral sequence for the morphism $f : \mathcal{X}_U \rightarrow U$:

$$H^p(U, R^q f_* \mathbf{Q}_l(n)) \implies H^{p+q}(\mathcal{X}_U, \mathbf{Q}_l(n)).$$

Here we denote by $R^q f_* \mathbf{Q}_l(n)$ the l -adic sheaf obtained from taking the higher direct images with finite coefficients. Given z we can extend it to a cycle on $\mathcal{X}_U$ of codimension n which is of degree zero on each fibre. This gives a cohomology class $\mathcal{Z}$ in

$$F_1 = \text{Ker}[H^{2n}(\mathcal{X}_U, \mathbf{Q}_l(n)) \rightarrow H^0(U, R^{2n} f_* \mathbf{Q}_l(n))].$$

From the spectral sequence there is a map:

$$F_1 \rightarrow H^1(U, R^{2n-1}f_*\mathbf{Q}_l(n)).$$

Claim: $\mathcal{Z}$ is zero in $H^1(U, R^{2n-1}f_*\mathbf{Q}_l(n))$. To prove the claim, we note that since z is in the Albanese kernel, its image in $H^1(k, H^{2n-1}(\overline{X}, \mathbf{Q}_l(n)))$ is zero. Then our claim follows from the following lemma:

Lemma 3.7: The map:

$$H^1(U, R^{2n-1}f_*\mathbf{Q}_l(n)) \rightarrow H^1(k, H^{2n-1}(\overline{X}, \mathbf{Q}_l(n)))$$

is injective.

Proof: By the smooth and proper base change theorem, the l -adic sheaf $\mathcal{F} = R^{2n-1}f_*\mathbf{Q}_l(n)$ is locally constant on U . Hence $H_v^1(U, \mathcal{F})=0$ for any closed point v of U and this gives the injectivity of the map.

From the spectral sequence and the vanishing of the cohomology class $\mathcal{Z}$ in $H^1(U, R^{2n-1}f_*\mathbf{Q}_l(n))$, we see that this class comes from $H^2(U, R^{2n-2}f_*\mathbf{Q}_l(n)) = H^2(G_S, H^{2n-2}(\overline{X}, \mathbf{Q}_l(n)))$. Our task is to show that this cohomology class vanishes in $H^2(k_v, H^{2n-2}(\overline{X}, \mathbf{Q}_l(n)))$ for all v in S .

Let $\mathcal{X}_v$ be a regular proper model of X_v over $\mathcal{O}_v$. By Proposition 3.2, after a finite extension of L/k_v , we can extend our original cycle z on X_v to an element of $CH^n(\mathcal{X}_v) \otimes \mathbf{Q}_l$ in such a way that its cohomology class lies in

$$H^{2n}(\mathcal{X}_v, \mathbf{Q}_l(n))^0 = \text{Ker}[H^{2n}(\mathcal{X}_v, \mathbf{Q}_l(n)) \rightarrow H^0(\mathcal{O}_v, R^{2n}f_*\mathbf{Q}_l(n))].$$

Since the natural map:

$$H^2(k_v, M) \rightarrow H^2(L, M)$$

is injective for any $\mathbf{Q}_l$ -vector space M , it suffices to prove that our cohomology class vanishes in $H^2(L, M)$.

Lemma 3.8:

$$H^{2n}(\mathcal{X}_v, \mathbf{Q}_l(n))^0 = 0.$$

This will prove the proposition.

Proof: From the Leray spectral sequence for $f : \mathcal{X}_v \rightarrow \text{Spec } \mathcal{O}_v$, we get the exact sequence:

$$0 \rightarrow H^2(\mathcal{O}_v, R^{2n-2}f_*\mathbf{Q}_l(n)) \rightarrow H^{2n}(\mathcal{X}_v, \mathbf{Q}_l(n))^0 \rightarrow H^1(\mathcal{O}_v, R^{2n-1}f_*\mathbf{Q}_l(n)) \rightarrow 0.$$

We claim that all three terms of this exact sequence are zero. The group on the left is zero because $\mathcal{O}_v$ has cohomological dimension one for torsion modules, hence for l -adic sheaves. As for the group on the right, two applications of the proper base change theorem give that it is isomorphic to $H^1(\mathbf{F}, H^{2n-1}(\overline{Y}, \mathbf{Q}_l(n)))$, where $\overline{Y}$ is the geometric special fibre of $\mathcal{X}_v$. By Deligne's theorem ([D2], 3.3.1), the geometric Frobenius acts on $H^{2n-1}(\overline{Y}, \mathbf{Q}_l(n))$ with weights which are ≤ -1 , hence nonzero. This shows that the group on the right vanishes and proves the lemma.

4 A Hasse Principle and Completion of the Proof

In this section we prove a ‘‘Hasse’’ principle in the function field case. In the case where Frobenius acts semi-simply on $H^{2n-2}(\overline{X}, \mathbf{Q}_l(n))$, this is proven in Jannsen's paper ([J2], Theorem 4 and Remark 7). The more general result given here was shown to me by Jannsen, and I am grateful to him for allowing me to give his proof in this paper. To avoid copying his earlier proof, we assume some familiarity with his argument.

Theorem 4.1 (Jannsen): Let k be a function field in one variable over a finite field $\mathbf{F}$ of characteristic p and l a prime number different from p . Let V be a $\mathbf{Q}_l$ -linear Galois representation of $G = \text{Gal}(k_{\text{sep}}/k)$ which is pure of weight -2 , and let S be a finite nonempty set of places of k including those for which the restriction to the decomposition group G_v is ramified. Then the natural map:

$$H^2(G_S, V) \rightarrow \bigoplus_{v \in S} H^2(G_v, V)$$

is injective.

Proof: Let Y be a smooth, projective model of k and let $U = Y - S, \overline{U} = U \times_{\mathbf{F}} \overline{\mathbf{F}}$ and $j : U \rightarrow Y$ the inclusion. Then by a theorem of Deligne ([D2], 3.4.1(iii)), V is a semi-simple $\pi_1(\overline{U})$ -representation. Then we can write $V = V_1 \oplus V_2$, where $\pi_1(\overline{U})$ acts trivially on V_2 and $V_1^{\pi_1(\overline{U})} = V_{1\pi_1(\overline{U})} = 0$. By Jannsen's proof in [J2], the result holds for V_1 , so we prove it for V_2 . Let F be the smooth $\mathbf{Q}_l$ -sheaf corresponding to V_2 . Then we have the exact sequence of cohomology groups:

$$(*) \dots H_S^2(Y, j_*F) \xrightarrow{g} H^2(Y, j_*F) \rightarrow H^2(U, F) \xrightarrow{f} H_S^3(Y, j_*F) \rightarrow \dots$$

By excision and the fact that $H^2(\mathcal{O}_v, j_*F) = H^2(\mathbf{F}, (j_*F)_{\mathbf{F}}) = 0$, we have:

$$H^2(k_v, F) = H_v^3(\mathcal{O}_v, j_*F)$$

for all $v \in S$, and the map f is the localization map. Thus an element of the kernel of the localization map comes from $H^2(Y, j_*F)$. It will then suffice to show that the map g in the exact sequence $(*)$ above is surjective.

Let $\Gamma = \text{Gal}(\overline{\mathbf{F}}/\mathbf{F})$ and consider the exact sequence of Γ -modules:

$$0 \rightarrow H^1(\Gamma, H^1(\overline{Y}, j_*F)) \rightarrow H^2(Y, j_*F) \rightarrow H^2(\overline{Y}, j_*F)^{\Gamma} \rightarrow 0.$$

By the fundamental theorem of Deligne ([D2], 3.2.3), the Γ -module $H^1(\overline{Y}, j_*F)$ is pure of weight -1, hence the left group is zero. Consider the following diagram:

$$\begin{array}{ccc} H_S^2(Y, j_*F) & \rightarrow & H_S^2(\overline{Y}, j_*F)^{\Gamma} \\ \downarrow & & \downarrow \\ H^2(Y, j_*F) & \rightarrow & H^2(\overline{Y}, j_*F)^{\Gamma} \end{array}.$$

We have just shown that the bottom horizontal map is an isomorphism and our goal is to show that the left vertical arrow is surjective. Since the top horizontal map is surjective, it will be enough to prove the surjectivity of the right vertical map. Let $\tilde{\mathcal{O}}_v$ denote the strict henselization of $\mathcal{O}_v$ and $\tilde{k}_v$ its fraction field. Then we have:

$$H^1(\tilde{k}_v, F) \cong H_v^2(\tilde{\mathcal{O}}_v, j_*F).$$

Now the pro- l -part of the absolute Galois group of $\tilde{k}$ is isomorphic to $\mathbf{Z}_l(1)$ as a Γ -module. Then by Pontryagin duality and the fact that $\pi_1(\overline{U})$ is assumed to act trivially on V_2 , we have

$$H^1(\tilde{k}_v, F)^\Gamma = (\hat{V}_2(1)_\Gamma)^*,$$

where $*$ denotes $\mathbf{Q}_l$ -linear dual and $\hat{V}_2$ denotes the dual representation. But by Poincaré duality, we have:

$$H^2(\overline{Y}, j_* F)^\Gamma \cong (\hat{V}_2(1)_\Gamma)^*,$$

and since S is nonempty, we obviously have injectivity of the dual map:

$$(\hat{V}_2(1)_\Gamma) \rightarrow \left[\bigoplus_{v \in S} \hat{V}_2(1)_\Gamma \right].$$

This completes the proof of the Theorem.

Now we can prove the main theorem of this paper:

Theorem 4.2: Let X be a smooth, projective, geometrically connected variety of dimension n over a function field k in one variable over a finite field of characteristic p , and l a prime number different from p . Assume Hypothesis 3.1 above. Then the map d_n^2 of section 1 is zero.

Proof: Combine Proposition 3.6 and Theorem 4.1, taking $V = H^{2n-2}(\overline{X}, \mathbf{Q}_l(n))$.

Theorem 4.3: Let X be a smooth, projective, geometrically connected surface over a function field in one variable over a finite field of characteristic $p \geq 5$. Then the map d_2^2 is zero.

Proof: Abhyankar [A] has proved resolution of singularities for 3-folds over an algebraically closed field of characteristic greater than or equal to 5. Given X , we can pass to a finite extension over which it has a smooth proper model. Then the kernel of d_2^2 will be killed by the degree of this extension, hence it is zero.

Appendix

In this appendix we prove a technical lemma which shows that the Abel-Jacobi map as defined via the Hochschild-Serre spectral sequence agrees with the map defined by Kummer theory on $Alb(X)$. This fact is probably known to several people, but I have been unable to find a reference.

Lemma: Let X be a smooth projective geometrically connected variety of dimension n over a field k , l a prime number different from $\text{char}.k$. Consider the map d_n^1 defined in §1 above and the map:

$$f_n^1 : A_0(X) \rightarrow H^1(k, H^{2n-1}(\overline{X}, \mathbf{Q}_l(n)))$$

defined as the composite of the Albanese map

$$\alpha : A_0(X) \rightarrow Alb(X)$$

and the map

$$Alb(X) \rightarrow H^1(k, H^{2n-1}(\overline{X}, \mathbf{Q}_l(n)))$$

obtained from Kummer theory on $Alb(X)$. Then $d_n^1 = f_n^1$.

Proof: First assume that X is a curve. The group $A_0(X)$ is generated by elements of the form $\text{cor}_{L/k}(P - Q)$, where L is a finite extension of k , P and Q are rational points of X_L and

$$\text{cor} : A_0(X_L) \rightarrow A_0(X)$$

denotes corestriction. Then it suffices to prove that for any such L/k and any two L -rational points P, Q of X , the two maps above give the same element in $H^1(L, H^1(\overline{X}, \mathbf{Z}_l(1)))$. We suppress L in what follows. Let $z = P - Q$, let Z be its support and $U = X - Z$. Taking cohomology with support in $\overline{Z}$, we easily get the exact sequence:

$$(*) : 0 \rightarrow H^1(\overline{X}, \mathbf{Z}/l^m(1)) \rightarrow H^1(\overline{U}, \mathbf{Z}/l^m(1)) \rightarrow \mathbf{Z}/l^m \rightarrow 0,$$

where the last group is generated by the class of z . Jannsen has shown ([J5], Lemma 9.4) that the extension given by $(*)$ gives the element of $H^1(k, H^1(\overline{X}, \mathbf{Z}/l^m(1)))$ which is associated to $P - Q$ via the map d_1^1 . Now consider the Kummer sequence for $Pic^0(\overline{X})$:

$$0 \rightarrow H^1(\overline{X}, \mathbf{Z}/l^m(1)) \rightarrow \text{Pic}^0(\overline{X}) \xrightarrow{l^m} \text{Pic}^0(\overline{X}) \rightarrow 0,$$

where the group on the left is identified with the l^m -torsion of $\text{Pic}^0(\overline{X})$. As is well-known, the boundary map:

$$\text{Pic}^0(\overline{X})^G \rightarrow H^1(k, H^1(\overline{X}, \mathbf{Z}/l^m(1)))$$

in the G -cohomology of this sequence may be obtained by pulling back the Kummer sequence by the G -map

$$\mathbf{Z} \rightarrow \text{Pic}^0(\overline{X})$$

corresponding to an element y of $\text{Pic}^0(\overline{X})^G$. One easily sees that the pullback extension is:

$$(**) : 0 \rightarrow H^1(\overline{X}, \mathbf{Z}/l^m(1)) \rightarrow D \rightarrow \mathbf{Z} \rightarrow 0,$$

where $D = \{x \in \text{Pic}^0(\overline{X}) : l^m x = ny \text{ for some } n \in \mathbf{Z}\}$. We may define a G -map

$$D/l^m \rightarrow H^1(\overline{U}, \mathbf{Z}/l^m(1))$$

as follows: Let $x \in D$, let $\mathcal{L}$ be the isomorphism class of line bundle associated to x and $\mathcal{M}$ the isomorphism class of line bundle associated to y . Choose an isomorphism

$$\mathcal{L}^{\otimes l^m} \rightarrow \mathcal{M}^{\otimes n}.$$

Such an isomorphism is unique modulo l^m , since if we had two such, then we would get an isomorphism from $\mathcal{O}_{\overline{X}}$ to itself. Since $\overline{X}$ is proper, this is given by an element of $\overline{k}^*$, and hence must be the identity modulo l^m . But this means that the two isomorphisms were the same modulo l^m . Since the class of y is supported on $\overline{Z}$, the restriction of this isomorphism to $\overline{U}$ gives a trivialization:

$$\mathcal{L}^{\otimes l^m} \cong \mathcal{O}_{\overline{U}},$$

and this data defines in a canonical way an element of $H^1(\overline{U}, \mathbf{Z}/l^m(1))$. It is easy to verify that via this map, the two extensions (*) and (**) give the same element of $H^1(k, H^1(\overline{X}, \mathbf{Z}/l^m(1)))$. Note that for this argument, we have used $\mathbf{Z}_l$ -coefficients.

In the case of X of arbitrary dimension, Poincaré duality and the Weil pairing give an identification:

$$H^{2n-1}(\overline{X}, \mathbf{Q}_l(n)) = V_l(\text{Alb}(\overline{X})).$$

Now given a zero-cycle z , if k is perfect, we can find a smooth, projective curve C going through it ([AK], [C]). By functoriality properties of the maps in question, we can easily reduce to the case of C which we have done above. If k is not perfect, such a curve C may be found after an inseparable extension of k . Then an easy restriction-corestriction argument shows that the maps are the same. This completes the proof of the lemma.

References

- [A] Abhyankar, S.: **Resolution of Singularities of Embedded Algebraic Surfaces**, Academic Press 1966
- [AK] Altman, A. and Kleiman, S. *Bertini theorems for hypersurface sections containing a subscheme*, Comm. in Algebra **7** (1979), 775-790
- [Be1] Beauville, A. *Quelques remarques sur la transformation de Fourier dans l'anneau de Chow d'une variété abélienne*, in **Algebraic Geometry: Tokyo/Kyoto 1982**, Lecture Notes in Mathematics 1016, Springer 1983
- [Be2] Beauville, A. *Sur l'anneau de Chow d'une variété abélienne*, Math. Ann. **273** (1986), 647-651
- [B1] Beilinson, A. *Height pairing on algebraic cycles*, in **Current Trends in Arithmetical Algebraic Geometry**, K. Ribet editor, Contemporary Mathematics Volume 67, American Mathematical Society 1987
- [B2] Beilinson, A. *Absolute Hodge cohomology*, in **Applications of Algebraic K-Theory to Algebraic Geometry and Number Theory**, S. Bloch, R.K. Dennis, E.M. Friedlander and M. Stein editors, Contemporary Mathematics Volume 55, American Mathematical Society 1986
- [Bl1] Bloch, S. *Algebraic cycles and values of L -functions*, J. für die Reine und Ang. Math. **350** (1984), 94-108
- [Bl2] Bloch, S. *K_2 of Artinian $\mathbf{Q}$ -algebras with application to algebraic cycles*, Communications in Algebra **3** (1975), 405-428
- [Bl3] Bloch, S. *Some elementary theorems about algebraic cycles on abelian varieties*, Inventiones Math. **37** (1976), 215-228
- [Bl4] Bloch, S. *Lectures on Algebraic Cycles*, Duke Univ. Press, 1980
- [BK] Bloch, S. and Kato, K. *L -functions and Tamagawa numbers of motives*, in **Grothendieck Festschrift**, Volume I, P. Cartier editor, Progress

in Mathematics, Birkhäuser 1990

[BLR] Bosch, S., Lütkebohmert, W. and Raynaud, M. **Néron Models**, Springer 1990

[CT-O] Colliot-Thélène, J.-L. and Ojanguren, M. *Variétés unirationnelles nonrationnelles: au delà de l'exemple d'Artin-Mumford*, Inventiones Math. **97** (1989), 141-158

[CT-R] Colliot-Thélène, J.-L. and Raskind, W. *Groupe de Chow de codimension deux des variétés définies sur un corps de nombres: un théorème de finitude pour la torsion*, Inventiones Math. **105** (1991), 221-245

[C] Coray, D. *Two remarks on the Bertini theorems*, unpublished typescript, 1980

[D1] Deligne, P. *Théorème de Lefschetz difficile et critères de dégénérescence de suites spectrales* Publ. Math. I.H.E.S. **35** (1968), 107-126

[D2] Deligne, P. *La Conjecture de Weil II*, Publ. Math. I.H.E.S. **52** (1980), 137-252

[E] Ekedahl, T. *On the adic formalism*, in **Grothendieck Festschrift**, Volume II, P. Cartier editor, Progress in Mathematics, Birkhäuser 1990

[Gi] Gillet, H. *Gersten's conjecture for the K-theory with torsion coefficients of a discrete valuation ring*, J. of Algebra **103** (1986), 377-380

[J1] Jannsen, U. *Continuous étale cohomology*, Math. Annalen **280** (1988), 207-245

[J2] Jannsen, U. *On the l -adic cohomology of varieties over number fields and its Galois cohomology*, in **Galois Groups over $\mathbb{Q}$** , Y. Ihara, K. Ribet and J.-P. Serre editors, Springer 1989

[J3] Jannsen, U. *letter to B. Gross*, 1989

- [J4] Jannsen, U. *Motivic sheaves and filtrations on Chow groups*, in **Motives: Seattle 1991**, U. Jannsen, S. Kleiman editors
- [J5] Jannsen, U. **Mixed Motives and Algebraic K-Theory**, Lecture Notes in Mathematics 1400, Springer 1990
- [L] Lang, S. **Diophantine Geometry**, Springer 1983
- [M1] Milne J.S. **Etale Cohomology**, Princeton Mathematical Series Volume 33, Princeton University Press 1980
- [M2] Milne, J.S. *Zero-cycles on algebraic varieties in non-zero characteristic: Rojzman's theorem*, Composition Math. **47** (1982), 271-287
- [Mu] Murre, J. *On a conjectural filtration on the Chow groups of an algebraic variety*, preprint 1992
- [N] Nori, M. *Algebraic cycles and Hodge-theoretic connectivity*, Inventiones Math. **105** (1993) 349-373
- [Ro] Roitman, A.A. *The torsion of the group of 0-cycles modulo rational equivalence*, Annals of Math. **111** (1980), 553-569
- [S1] Saito, S. *Motives and filtrations on Chow groups*, preprint 1993
- [S2] Saito, S. *On the cycle map for torsion algebraic cycles of codimension two*, Inventiones Math. **106** (1991), 443-460
- [S3] Saito, S. *Remarks on algebraic cycles*, preprint 1993
- [SH] Saito, H. *Generalization of Abel's Theorem and some finiteness properties of zero-cycles on surfaces*, preprint
- [T] Thomason, R. *Absolute Cohomological Purity*, Bull. Math. Soc. France **112** (1984), 397-406